

Short-range stationary patterns and long-range disorder in an evolution equation for one-dimensional interfaces

J. Munoz-Garcia, R. Cuerno, M. Castro

Abstract- A local evolution equation for one-dimensional interfaces is derived in the context of erosion by ion beam sputtering. We present numerical simulations of this equation which show interrupted coarsening in which an ordered cell pattern develops with constant wavelength and amplitude at intermediate distances, while the profile is disordered and rough at larger distances. Moreover, for a wide range of parameters the lateral extent of ordered domains ranges up to tens of cells. We also provide analytical estimates for the stationary pattern wavelength and mean growth velocity.

Index Terms- ion-sputtered surfaces, aeolian sand ripples, thin-film growth, instabilities, model

Due to copyright restriction we cannot distribute this content on the web. However, clicking on the next link, authors will be able to distribute to you the full version of the paper:

[Request full paper to the authors](#)

If your institution has an electronic subscription to Physical Review E, you can download the paper from the journal website:

[Access to the Journal website](#)

Citation:

Munoz-Garcia, J.; Cuerno, R.; Castro, M.; "Short-range stationary patterns and long-range disorder in an evolution equation for one-dimensional interfaces", Physical Review E, vol.74, no.5, pp.050103.1-050103.4. November, 2006.