

# Plasma distributions observed in a 2.45 GHz hydrogen discharge

O.D. Cortázar; A.M. Megia Macías; O. Tarvainen; A. Vizcaíno de Julián

## Abstract-

The existence of various spatial distributions of hydrogen plasma in a pulsed 2.45 GHz microwave discharge is demonstrated. The data has been obtained through optical emission diagnostics utilizing an ultra-fast CCD camera system with multi-channel plate (MCP) intensifiers, and a wavelength-filtered photodiode recording temporal light emission signals of hydrogen atoms and molecules. It has been observed that the magnetic field topology and strength are determining the transitions between different plasma patterns and spectral saturation times while neutral gas pressure and microwave power show a weaker influence on the profiles but affect the emitted light intensity.

## Index Terms-

Due to copyright restriction we cannot distribute this content on the web. However, clicking on the next link, authors will be able to distribute to you the full version of the paper:

[Request full paper to the authors](#)

If your institution has an electronic subscription to Plasma Sources Science and Technology, you can download the paper from the journal website:

[Access to the Journal website](#)

## Citation:

*Cortázar, O.D.; Megia-Macías, A.; Tarvainen, O.; Vizcaíno-de-Julián, A. "Plasma distributions observed in a 2.45 GHz hydrogen discharge", Plasma Sources Science and Technology, vol.23, no.6, pp.065028-1-065028-12, December, 2014.*